

Dr. med. Kathrin Liebmann

## Contact details

### Practice Berlin-Kladow

Sakrower Landstr. 29

14089 Berlin

Tel: 030.364 330 88/87

Opening times: Mon, Wed, Fri 9–14 | Tue, Thu 14–19

All insured and private patients

### Private practice in Potsdam

Alleestr. 13

14469 Potsdam

Tel: 0331. 201 57 00

Appointments by prior arrangement.

Private patients and “Beihilfe” arrangements.

Email: [info@dr-liebmann.de](mailto:info@dr-liebmann.de)

[www.dr-liebmann.de](http://www.dr-liebmann.de)



Antenatal care  
and obstetrics

# Dear patient!

Welcome to our obstetric practice and thank you for choosing our modern and service-oriented team.

Our practice philosophy is to provide you with the latest and most up-to-date antenatal care, with an emphasis on clear and thorough consultation.

You will find our team members friendly and helpful, and good appointment planning ensures that waiting times at our practices are generally kept below 30 minutes.

## Services

In addition to the standard antenatal screening services covered by the statutory health insurance in Germany, we recommend the following screening options, which are tried and tested methods for ensuring that any complications during pregnancy are detected early:

- Toxoplasmosis screening
- Nuchal translucency scan
- Glucose tolerance test
- Additional ultrasound scans with placental bloodflow measurement
- Streptococcus screening

These screens are covered by private health insurance policies. However, unfortunately, they are not covered by the statutory health insurance and will have to be paid for privately as so-called Individual Healthcare Services (known within the German healthcare system under the acronym “IGeL”).

We offer these “IGeL” services for the following reasons: The proportion of pregnant women affected by toxoplasmosis infection is 0.7%, gestational diabetes (pregnancy diabetes) affects 3–5% of pregnant women, placental insufficiency occurs in 4% of pregnancies, and streptococcus infections affect 16% of pregnant women. Early detection of complications during pregnancy ensures that appropriate therapeutic and preventive steps can be taken early on. The chance of having a baby with Down’s syndrome (trisomy 21) is 0.3% for a 35-year-old, 1% for a 40-year-old mother.

We believe that the only sensible preventive medical care lies in using the best and most up-to-date screening technology. We therefore place great emphasis on keeping abreast of the latest developments and technological advances.

Where required, we will be happy to arrange for further consultation through colleagues with specific expertise, including our midwife, or refer you to hospitals with which we have close links and where you will be well looked after.